

# **The Canadian Entomologist**

**INDEX TO VOLUME 109  
1977**

**Editor: P. E. Morrison**



**THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA**

- Akre, R.D. See Novak, J. A., 663.
- Alauzet, C. Cycle biologique de *Pissodes notatus* (Coleoptera: Curculionidae) dans la région Toulousaine (France), 597.
- AliNiazee, M.T. and Brown, R.D. Laboratory rearing of the western cherry fruit fly, *Rhagoletis indifferens* (Diptera: Tephritidae): Oviposition and larval diets, 1227.
- Allen, R.K. and Cohen, S.D. Mayflies (Ephemeroptera) of Mexico and Central America: New species, descriptions, and records, 399.
- Alleyne, E.H. and Morrison, F.O. Some Canadian poplar aphid galls, 321.
- Allyson, S. A study of some North American larvae of the genus *Tetralopha* Zeller (Lepidoptera: Pyralidae: Epipaschiinae), 329.
- The larvae of two species of Macrothecinae (Lepidoptera: Pyralidae), 839.
- Arthur, A.P. and Kell, W.L. A mechanical method of mass collecting lepidopterous pupae and parasite cocoons from infested soil, 895.
- Atwood, C.E. See Hartwick, E.B., 129.
- Ayre, G.L. See Struble, D.L., 975.
- Bailey, B.K. See Steck, W., 157.
- Bailey, C. G. and Mukerji, M.K. Energy dynamics of *Melanoplus bivittatus* and *M. femurrubrum* (Orthoptera: Acrididae) in a grassland ecosystem, 605.
- Bailey, C.G. and Singh, N.B. An energy budget for *Mamestra configurata* (Lepidoptera: Noctuidae), 687.
- Baker, B.H., Hostettler, B.B., and Furniss, M.M. Response of eastern larch beetle (Coleoptera: Scolytidae) in Alaska to its natural attractant and to Douglas-fir beetle pheromones, 289.
- Baldwin, W.F. See Nelson, D.L., 319.
- Baldwin, W.F., Gross, H.P., Wilson, M.L., Keill, D.J., Stuart, R.J., Sebastien, R.J., Knight, A.G., Chant, G.D., Knight, P.A., and West, A.S. Suppression of black fly populations in Deep River, Ontario, 249.
- Barbosa, P. See Capinera, J.L., 1313.
- Barclay, H. and van den Driessche, P. Predator-prey models with added mortality, 763.
- Barlow, C. A., Randolph, P.A., Randolph, J.C. Effects of pea aphids, *Acyrtosiphon pisum* (Homoptera: Aphididae), on growth and productivity of pea plants, *Pisum sativum*, 1491.
- Barfield, C.S., Sharpe, P.J.H., and Bottrell, D.G. A temperature-driven developmental model for the parasite *Bracon mellitor* (Hymenoptera: Braconidae), 1503.
- Barnard, D.R. and Mulla, M.S. Temporal patterns of pupation and emergence in the mosquito *Culiseta inornata* (Diptera: Culicidae), 1387.
- Batth, S.S. A survey of Canadian populations of the German cockroach for resistance to insecticides, 49.
- Bélanger, A. See Martel, P., 317.
- Bergeron, J.-M. See Robert, S., 1571.
- Berryman, A.A. See Bordsach, R.P., 95.
- Beveridge, W.J.G. See Kingsbury, P.D., 1265.
- Binns, M.R. See Harcourt, D.G., 1521.
- Bodnaryk, R.P. A melanotic mutant of the moth *Mamestra configurata* (Lepidoptera: Noctuidae), 479.
- Boiteau, G. and Perron, J.-M. Etude du complexe hémocytaire postembryonnaire de *Lambdina fiscellaria fiscellaria* (Lepidoptera: Geometridae), 269.
- Bolte, K.B. A new species of *Eupithecia* (Lepidoptera: Geometridae) from Alaska, 1019.
- See also Morris, Ray F., 385.
- Bond, E.J. and Svec, H.J. Control of Colorado potato beetle, *Leptinotarsa decemlineata*, in shipments of harvested potatoes, 285.
- Bordsach, R.P. and Berryman, A.A. Host resistance to the fir engraver beetle, *Scolytus ventralis* (Coleoptera: Scolytidae). 2. Repellency of *Abies grandis* resins and some monoterpenes, 95.
- Borden, J.H. See McLean, J.A., 675.
- Borg, T.K. See Stevens, R.E., 1269.
- Bottrell, D.G. See Barfield, C.S., 1503.
- Bracken, G.K. See Bucher, G.E., 549; Wylie, H.G., 1295.
- Brewer, J.W. See Yarger, L.C., 849.
- Brewer, J.W. and O'Neal, J. Early ground application of acephate for control of the western spruce budworm (Lepidoptera: Tortricidae) on Douglas-fir, 1153.
- Bright, D.E. New synonymy, new combinations, and new species of North American *Pityophthorus* (Coleoptera: Scolytidae). I, 511.
- Brown, E.A. and Cameron, E.A. Studies of the compound eye of *Lymantria dispar* (Lepidoptera: Lymantriidae) males, and behavioral implications, 255.
- Brown, R.D. See AliNiazee, M.T., 1227.
- Bryan, J.M. and Elvidge, J. Mortality of adult grain beetles in sample delivery systems used in terminal grain elevators, 209.

- Bryant, D.G.** See Raske, A.G., 1297, 1307.
- Bucher, G.E.** See Wylie, H.G., 823.
- Bucher, G.E. and Bracken, G.K.** The bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae). Sterilization of adults by exposure of prepupal-pupal stages to 25°C, 549.
- Buffam, P.E.** See Frye, R.H., 1221.
- Burgess, L.** Flea beetles (Coleoptera: Chrysomelidae) attacking rape crops in the Canadian Prairie Provinces, 21.
- Geocoris bullatus*, an occasional predator of flea beetles (Hemiptera: Lygaeidae), 1519.
- Burrage, R.H.** See McKinlay, K.S., 871.
- Butler, G.D.** See Gutierrez, A.P., 1457.
- Cadogan, B.L. and Laing, J.E.** A technique for rearing the predaceous mite *Balaustium putmani* (Acarina: Erythraeidae) with notes on its biology and life history, 1535.
- Cameron, E.A.** See Brown, E.A., 255.
- Cameron, P.J. and Morrison, F.O.** Analysis of mortality in the apple maggot, *Rhagoletis pomonella* (Diptera: Tephritidae), in Quebec, 769.
- Campbell, A. and Mackauer, M.** Reproduction and population growth of the pea aphid (Homoptera: Aphididae) under laboratory and field conditions, 277.
- Capinera, J.L. and Barbosa, P.** Influence of natural diets and larval density on gypsy moth, *Lymantria dispar* (Lepidoptera: Orgyiidae), egg mass characteristics, 1313.
- Carolin, V.M.** See Medley, R.D., 1121.
- Caron, R.E.** See Stinner, R.E., 879.
- Cave, R.D.** See Gutierrez, A.P., 1375.
- Chant, D.A., Hansell, R.I.C., and Rowell, H.J.** An analysis of interspecific and intraspecific morphological differences in two closely related species of *Amblyseius* Berlese using methods of numerical taxonomy (Acarina: Phytoseiidae), 1605.
- Chant, G.D.** See Baldwin, W.F., 249.
- Chawla, S.S.** See Perusse, F., 267.
- Cheng, H.H.** Insect parasites of the darksided cutworm, *Euxoa messoria* (Lepidoptera: Noctuidae), in Ontario, 137.
- Flue-cured tobacco losses caused by the tomato hornworm, *Manduca quinquemaculata* (Lepidoptera: Sphingidae), at various infestation levels in Ontario, 1091.
- Chiang, H.C.** See Okubo, A., 149.
- Chiang, M.S.** See Hudon, M., 931.
- Chillcott, J.C.** See Teskey, H.J., 1445.
- Chisholm, M.D.** See Steck, W., 157; Underhill, E.W., 1335; Struble, D.L., 1393.
- Christensen, J.B., Gutierrez, A.P., Cothran, W.R., and Summers, C.G.** The within field spatial pattern of the larval Egyptian alfalfa weevil, *Hypera brunneipennis* (Coleoptera: Curculionidae): An application of parameter estimates in simulation, 1599.
- Church, N.S. and Gerber, G.H.** The development and habits of *Linsleya sphaericollis* (Coleoptera: Meloidae), 375.
- Observations on the ontogeny and habits of *Lytta nuttalli*, *L. viridana*, and *L. cyanipennis* (Coleoptera: Meloidae): The adults and eggs, 565.
- Cohen, S.D.** See Allen, R.K., 399.
- Collins, N.C.** Mechanisms determining the relative abundance of brine flies (Diptera: Ephydriidae) in Yellowstone thermal spring effluents, 415.
- Copper, W.A.** See Dahlsten, D.L., 727.
- Cothran, W.R.** See Christensen, J.B., 1599.
- Coulson, R.N.** See Pulley, P.E., 39; 1325.
- Coulson, R.N., Pulley, P.E., Foltz, J.L., Martin, W.C., and Kelley, C.L.** Survival models for within-tree populations of *Dendroctonus frontalis* (Coleoptera: Scolytidae), 1701.
- Craig, G.B.** See Ramalingam, S., 897.
- Croft, B.A. and Hoying, S.A.** Competitive displacement of *Panonychus ulmi* (Acarina: Tetranychidae) by *Aculus schlechtendali* (Acarina: Eriophyidae) in apple orchards, 1025.
- Curry, G.L. and DeMichele, D.W.** Stochastic analysis for the description and synthesis of predator-prey systems, 1167.
- Dahlsten, D.L.** See Ohmart, C.P., 1001.
- Dahlsten, D.L., Luck, R.F., Schlinger, E.I., Wenz, J.M., and Copper, W.A.** Parasitoids and predators of the Douglas-fir tussock moth, *Orgyia pseudotsugata* (Lepidoptera: Lymantridae), in low to moderate populations in central California, 727.
- Dateman, G.E.** See Sanders, C.J., 1203.
- Dateman, G.E., Robbins, R.G., Eichlin, T.D., Pierce, J.** Forest Lepidoptera attracted by known sex attractants of western spruce budworms, *Choristoneura* spp. (Lepidoptera: Tortricidae), 875.
- Daxl, R.** See Wang, Y., 1359.
- Delisle, J.** See McNeil, J.N., 1199.

- DeMichele, D.W. See Curry, G.L., 1167.
- Devlin, C.G. See Steck, W., 157.
- Dew, H.E. See Gibo, D.L., 615.
- Doane, J.F. Spatial pattern and density of *Ctenicera destructor* and *Hypolithus bicolor* (Coleoptera: Elateridae) in soil in spring wheat, 807.
- Dowell, R.V. and Horn, D.J. Adaptive strategies of larval parasitoids of the alfalfa weevil (Coleoptera: Curculionidae), 641.
- Downing, R.S. and Logan, D.M. A new approach to San Jose scale control (Hemiptera: Diaspididae), 1249.
- Duchesne, R.-M. See McNeil, J.N., 1253.
- Dunphy, G.B., Keough, K.M.W., and Nolan, R.A. Fatty acid composition and lipid content of late larval and pupal stages of the eastern hemlock looper, *Lambdina fiscellaria fiscellaria* (Lepidoptera: Geometridae), 347.
- Dunphy, G.B., Nolan, R.A., and Otvos, I.S. Ninhydrin-positive substance analysis of larval hemolymph of the eastern hemlock looper, *Lambdina fiscellaria fiscellaria* (Lepidoptera: Geometridae) and growth of *Entomophthora egressa* protoplasts, 341.
- Duval, A. See Perusse, F., 267.
- Dyer, E.D.A. and Hall, P.M. Factors affecting larval diapause in *Dendroctonus rufipennis* (Coleoptera: Scolytidae), 1485.
- Dyer, E.D.A. and Safranyik, L. Assessment of the impact of pheromone-baited trees on a spruce beetle population (Coleoptera: Scolytidae), 77.
- Ebbesmeyer, C.C. See Okubo, A., 149.
- Eichlin, T.D. See Daterman, G.E., 875.
- Eikenbary, R.D., Grovenburg, W.G., Hedger, G.H., and Morrison, R.D. Modification and further evaluation of an equation for predicting populations of *Curculio caryae* (Coleoptera: Curculionidae), 1159.
- Elliott, W.M. Mating frequency of the female European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae), in southwestern Ontario, 117.
- Ellis, C.R. See Harcourt, D.G., 473.
- Elsey, K.D. Dissemination of *Howardula* sp. nematodes by adult tobacco flea beetles (Coleoptera: Chrysomelidae), 347.
- Elvidge, J. See Bryan, J.M., 209.
- Emory, R.W. See Reisen, W.K., 1475, 1481.
- Enfield, M.A. and Pritchard, G. Estimates of population size and survival of immature stages of four species of *Aedes* (Diptera: Culicidae) in a temporary pond, 1425.
- Methods for sampling immature stages of *Aedes* spp. (Diptera: Culicidae) in temporary ponds, 1435.
- Ennis, T.J. See Sanders, C.J., 1203.
- Fast, P.G. *Bacillus thuringiensis*  $\delta$ -endotoxin: On the relative roles of spores and crystals in toxicity to spruce budworm (Lepidoptera: Tortricidae), 1515.
- Fernando, C.H. See Galbraith, D.F., 221.
- Finnegan, R.J. Establishment of a predacious red wood ant, *Formica obscuripes* (Hymenoptera: Formicidae), from Manitoba to eastern Canada, 1145.
- See also McNeil, J.N., 1199.
- Fisher, R.W. and Menzies, D.R. A rapid, precise method of exposing oriental fruit moth (Lepidoptera: Olethreutidae) larvae to pesticide deposits, 1399.
- Fogal, W.H. Effect of a phenyl-benzoyl urea [1-(4-chlorophenyl)-3-(2,6-difluorobenzoyl)-urea] on *Diprion similis* (Hymenoptera: Diprionidae), 981.
- Foltz, J.L. See Pulley, P.E., 39, 1325; Coulson, R.N., 1071.
- Foott, W.H. Biology of the corn leaf aphid, *Rhopalosiphum maidis* (Homoptera: Aphididae), in southwestern Ontario, 1129.
- Friend, W.G. See Hartwick, E.B., 129; Smith, J.J.B., 477.
- Friend, W.G. and Smith, J.J.B. An apparatus for exchanging diets rapidly during artificial feeding of sucking insects, 15.
- Frye, R.H., Schmid, J.M., Lister, C.K., and Buffam, P.E. Post-attack injection of Silvisar 510 (cacodylic acid) in spruce beetle (Coleoptera: Scolytidae) infested trees, 1221.
- Furniss, M.M. See Baker, B.H., 289.
- Furniss, M.M., Young, J.W., McGregor, M.D., Livingston, R.L., and Hamel, D.R. Effectiveness of controlled-release formulations of MCH for preventing Douglas-fir beetle (Coleoptera: Scolytidae) infestation in felled trees, 1063.
- Gage, S.H. See Mukerji, M.K., 229.
- Galbraith, D.F. and Fernando, C.H. The life history of *Gerris remigis* (Heteroptera: Gerridae) in a small stream in southern Ontario, 221.
- Garnett, W.B. See Novak, J.D., 663.
- Gaudet, P.M. See Harvey, G.T., 987.

- Gerber, G.H.** See Church, N.S., 375, 565.
- Gibbs, K.E.** Evidence for obligatory parthenogenesis and its possible effect on the emergence period of *Cloeon triangulifer* (Ephemeroptera: Baetidae), 337.
- Gibo, D.L.** A method for rearing various species of social wasps of the genus *Polistes* (Hymenoptera: Vespidae) under controlled conditions, 1013.
- Gibo, D.L., Temporale, A., Lamarre, T.P., Soutar, B.M., and Dew, H.E.** Thermoregulation in colonies of *Vespula arenaria* and *V. maculata* (Hymenoptera: Vespidae). III. Heat production in queen nests, 615.
- Golini, V.I.** Simuliidae (Diptera) of Rendalen, Norway. III. Females of the *Eusimulium rendalense* complex infected with mermithids (Nematoda), 755.
- Granett, J. and Retnakaran, A.** Stadial susceptibility of eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), to the insect growth regulator Dimilin®, 893.
- Griffiths, K.J.** See Sullivan, C.R., 215.
- Gross, H.P.** See Baldwin, W.F., 249.
- Grovenburg, W.G.** See Eikenbary, R.D., 1159.
- Guevremont, H.C. and Quednau, F.W.** Morphologie et biologie de *Grypocentrus albipes* (Hymenoptera: Ichneumonidae), parasite de la petite mineuse du bouleau, *Fenusa pusilla* (Hymenoptera: Tenthredinidae), 1417.
- Introduction de parasites Ichneumonides pour la lutte biologique contre *Fenusa pusilla* (Hymenoptera: Tenthredinidae) au Québec, 1545.
- Guppy, J.C.** See Harcourt, D.G., 473, 1521.
- Guppy, J.C. and Harcourt, D.G.** Population assessment during the adult stage of the alfalfa weevil, *Hypera postica* (Coleoptera: Curculionidae), 497.
- Gutierrez, A.P.** See Wang, Y., 1359; Christensen, J.B., 1599.
- Gutierrez, A.P., Butler, G.D., Wang, Y., and Westphal, D.** The interaction of pink bollworm (Lepidoptera: Gelichiidae), cotton, and weather: A detailed model, 1457.
- Gutierrez, A.P., Leigh, T.F., Wang, Y., and Cave, R.D.** An analysis of cotton production in California: *Lygus hesperus* (Heteroptera: Miridae) injury — an evaluation, 1375.
- Gwynne, D.T.** Mating behavior of *Neoconocephalus ensiger* (Orthoptera: Tettigoniidae) with notes on the calling song, 237.
- Hagley, E.A.C.** Effect of low temperatures on mortality of hibernating codling moth larvae and fertility of surviving adults (Lepidoptera: Olethreutidae), 123.
- See also Jaques, R.P., 1079; Menzies D.G., 1405.
- Hall, P.M.** See Dyer, E.D.A., 1485.
- Halliday, C.G.** See Smith, B.C., 59.
- Hamel, D.R.** The effects of *Bacillus thuringiensis* on parasitoids of the western spruce budworm, *Choristoneura occidentalis* (Lepidoptera: Tortricidae), and the spruce coneworm, *Dioryctria reniculelloides* (Lepidoptera: Pyralidae), in Montana, 1409.
- See also Furniss, M.M., 1063.
- Hamilton, K.G.A.** Review of the World species of *Prosapia* Fennah (Rhynchota: Homoptera: Cercopidae), 621.
- Hamilton, W.J. and Penrith, M-L.** Description of an individual possible hybrid tenebrionid beetle and the habitat preference of the parental species, 701.
- Hansell, R.I.C.** See Chant, D.A., 1605.
- Harcourt, D.G.** See Guppy, J.C., 497.
- Harcourt, D.G., Guppy, J.C., and Binns, M.R.** The analysis of intrageneration change in eastern Ontario populations of the alfalfa weevil, *Hypera postica* (Coleoptera: Curculionidae), 1521.
- Harcourt, D.G., Guppy, J.C., and Ellis, C.R.** Establishment and spread of *Tetrastichus julis* (Homoptera: Eulophidae), a parasitoid of the cereal leaf beetle in Ontario, 473.
- Harris, C.R.** Biological activity of chlorpyrifos, chlorpyrifos-methyl, phorate, and Counter® in soil, 1115.
- Harris, C.R. and Turnbull, S.A.** Laboratory assessment of the potential of 16 experimental compounds as soil insecticides, 1109.
- Hartwick, E.B., Friend, W.G., and Atwood, C.E.** Trail-laying behaviour of the carpenter ant, *Camponotus pennsylvanicus* (Hymenoptera: Formicidae), 129.
- Harvey, G.T.** Mean weight and rearing performance of successive egg clusters of eastern spruce budworm (Lepidoptera: Tortricidae), 487.
- Harvey, G.T. and Gaudet, P.M.** The effects of benomyl on the incidence of microsporidia and the developmental performance of eastern spruce budworm (Lepidoptera: Tortricidae), 987.
- Hedger, G.H.** See Eikenbary, R.D., 1159.
- Holliday, N.J.** Sex determination in living adult ground beetles (Coleoptera: Carabidae), 397.
- Holling, C.S.** See Thomson, A.J., 1181.
- Holmes, N.D.** The effect of the wheat stem sawfly, *Cephus cinctus* (Hymenoptera: Cephidae), on the yield and quality of wheat, 1591.
- See also Smith, D.S., 575.
- Horn, D.J.** See Dowell, R.V., 641.

- Hostettler, B.B.** See Baker, B.H., 289.
- House, H.L.** Interaction between amino acids and glucose in larval nutrition of the fly *Agria housei* (Diptera: Sarcophagidae) at low temperature, 87.
- Hoying, S.A.** See Croft, B.A., 1025.
- Hudon, M. and Chiang, M.S.** Influence of resistant and susceptible maize inbred lines on the biology of the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae), after natural overwintering conditions in southwestern Quebec, 931.
- Hudson, A.** Additional records of Newfoundland Tabanidae, 639.
- Huggert, L.** Note on the occurrence of *Anopodias lacustris* in Canada (Hymenoptera: Proctotrupoidea: Platygasterinae), 107.
- Hughes, K.M.** Observations on the morphology of nuclear polyhedrosis viruses from six forest insects, 759.
- Ives, W.G.H.** Trends in populations of three species of sawflies infesting tamarack in southeastern Manitoba (Hymenoptera: Tenthredinidae), 635.
- James, M.T.** The South American species of *Oplachantha* (Diptera: Stratiomyidae), 305.
- Jakes, R.P., MacLellan, C.R., Sanford, K.H., Proverbs, M.D., and Hagley, E.A.C.** Preliminary orchard tests on control of codling moth larvae by a granulosis virus, 1079.
- Jay, S.C.** See Ying-Shin Peng, 1175.
- Jones, J.W.** See Stinner, R.E., 879.
- Jones, R.A.** See Rice, R.E., 1403.
- Jong, C.** See McMullen, R.D., 165.
- Kay, C.A.R., Veazey, J.N., and Whitcomb, W.H.** Effects of date of soil disturbance on numbers of adult field crickets (Orthoptera: Gryllidae), in Florida, 721.
- Kearby, W.H.** See Roling, M.P., 1235.
- Keill, D.J.** See Baldwin, W.F., 249.
- Kell, W.L.** See Arthur, A.P., 895.
- Kelley, C.L.** See Coulson, R.N., 1071.
- Kelton, L.A.** New species of *Cardiastethus* Fieber and *Melanocoris* Champion, and new records of European *Acomporis* Reuter and *Tennostethus* Fieber in Canada (Heteroptera: Anthocoridae), 243.  
A new species of *Elatophilus* Reuter from Ontario, and new synonymy for *Piezostethus flaccidus* Van Duzee (Heteroptera: Anthocoridae), 1017.  
Species of the genus *Pinalitus* Kelton found in Canada (Heteroptera: Miridae), 1549.
- Keough, K.M.W.** See Dunphy, G.B., 347.
- Kingsbury, P.D. and Beveridge, W.J.G.** A simple bubbler for sorting bottom fauna samples by elutriation, 1265.
- Kirtibutr, N. and Schenk, J.A.** Differentiation of larval *Pityokteines elegans* and *Scolytus ventralis* (Coleoptera: Scolytidae), 381.
- Knight, A.G.** See Baldwin, W.F., 249.
- Knight, P.A.** See Baldwin, W.F., 249.
- Laing, J.E.** See White, N.D., 1275; Cadogan, B.L., 1535.
- Lamarre, T.P.** See Gibo, D.L., 615.
- Latheef, M.A.** See Ng, Y.S., 1257.
- Latheef, M.A., Yeargan, K.V., and Pass, B.C.** Effect of density on host-parasite interactions between *Hypera postica* (Coleoptera: Curculionidae) and *Bathyplectes anurus* (Hymenoptera: Ichneumonidae), 1057.
- Leigh, T.F.** See Gutierrez, A.P., 1375.
- Letendre, M.** See McNeil, J.N., 37.
- Lilly, C.E.** See Struble, D.L., 261.
- Lister, C.K.** See Frye, R.H., 1221.
- Livingston, R.L.** See Furniss, M.M., 1063.
- Loeffler, J.** See Steck, W., 157.
- Logan, D.M.** See Downing, R.S., 1249.
- Luck, R.F.** See Dahlsten, D.L., 727.
- Lyons, L.A.** Parasitism of *Neodiprion sertifer* (Hymenoptera: Diprionidae) by *Exenterus* spp. (Hymenoptera: Ichneumonidae) in Ontario, 1962-1972, with notes on the parasites, 555.
- MacKauer, M.** See Campbell, A., 277.
- MacKay, P.A. and Wellington, W.G.** Notes on the life history and habits of the red-backed sawfly, *Eriocampa ovata* (Hymenoptera: Tenthredinidae), 53.
- MacLellan, C.R.** Populations of some major pests and their natural enemies on young and semidwarf apple trees in Nova Scotia, 797.  
Trends of codling moth (Lepidoptera: Olethreutidae) populations over 12 years on two cultivars in an insecticide free orchard, 1555.  
See also Jakes, R.P., 1079.
- Madder, M.C.A., Rosenberg, D.M., and Wiens, A.P.** Larval cocoons in *Eukiefferiella claripennis* (Diptera: Chironomidae), 891.

- Madsen, H.F., Potter, S.A., and Peters, F.E. Pest management: Control of *Archips argyrospilus* and *Archips rosanus* (Lepidoptera: Tortricidae) on apple, 171.
- Martel, P. and Bélanger, A. Xerography, a new tool for in situ study of insects, 317.
- Martin, W.C. See Pulley, P.E., 39, 1325; Coulson, R.N., 1071.
- Masner, L. A new genus of ecitophilous diapiiid wasps from Arizona (Hymenoptera: Proctotrupeidae: Diapriidae), 33.
- Massalski, A. See Philogène, B.J.R., 143.
- Mayyasi, A.M. See Pulley, P.E., 39.
- McEwen, F.L. See Reid, J.A.K., 1287.
- McGregor, M.D. See Furniss, M.M., 1063.
- McKinlay, K.S. and Burrage, R.H. Laboratory studies on the toxicities of 10 insecticides to the grasshopper, *Melanoplus sanguinipes* (Orthoptera: Acrididae). I, 871.
- McLean, J.A. and Borden, J.H. Attack by *Gnathotrichus sulcatus* (Coleoptera: Scolytidae) on stumps and felled trees baited with sulcatol and ethanol, 675.
- McLeod, J.M. Distribution of ovipositional attacks by parasitoids on overwintering larvae of the spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), 789.
- McMullen, R.D. and Jong, C. Effect of temperature on developmental rate and fecundity of the pear psylla, *Psylla pyricola* (Homoptera: Psyllidae), 165.
- McNamara, J. A catalogue of types of Coleoptera in the Canadian National Collection of Insects, Supplement I, 175.
- McNeil, J.N., Delisle, J., and Finnegan, R.J. Inventory of aphids on seven conifer species in association with the introduced red wood ant, *Formica lugubris* (Hymenoptera: Formicidae), 1199.
- McNeil, J.N. and Duchesne, R.-M. Transport of hay and its importance in the passive dispersal of the European skipper, *Thymelicus lineola* (Lepidoptera: Hesperidae), 1253.
- McNeil, J.N., Smirnov, W.A., and Letendre, M. *Bacillus thuringiensis* as a means of control for the European skipper, *Thymelicus lineola* (Lepidoptera: Hesperidae), 37.
- Medley, R.D. and Carollin, V.M. *Chaetophleps nasellensis*, a dipterous parasitoid of the western hemlock looper (Lepidoptera: Geometridae), 1121.
- Menzies, D.R. See Fisher, R.W., 1399.
- Menzies, D.R. and Hagley, E.A.C. A mechanical trap for sampling populations of small, active insects, 1405.
- Morris, O. N. Long-term effects of aerial applications of virus-fenitrothion combinations against the spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), 9.  
Compatibility of 27 chemical insecticides with *Bacillus thuringiensis* var. *kurstaki*, 855.  
Long term study of the effectiveness of aerial application of *Bacillus thuringiensis* - acephate combinations against the spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), 1239.  
Relationship between microbial numbers and droplet size in aerial spray application, 1319.
- Morris, Ray F. and Bolte, K. The European species *Perizoma alchemilla* (the small rivulet (Br.)) (Lepidoptera: Geometridae) in Newfoundland, 385.
- Morrison, F.O. See Alleyne, E.H., 321; Cameron, P.J., 769.
- Morrison, R.D. See Eikenbary, R.D., 1159.
- Mukerji, M.K., Gage, S.H., and Randell, R.L. Influence of embryonic development and heat on population trend of three grasshopper species (Orthoptera: Acrididae) in Saskatchewan, 229.
- Mukerji, M.K. See Bailey, C.G., 605.
- Mulla, M.S. See Barnard, D.R., 1387.
- Munroe, E. Two species of *Palpita* Hübner described by Janse, with descriptions of new species (Lepidoptera: Pyralidae: Pyraustinae), 67.  
Synopsis of *Heortia* Lederer, with descriptions of a new species and a new subspecies (Lepidoptera: Pyralidae: Odontiinae), 429.  
A new genus and two new species of Neotropical Pyraustinae (Lepidoptera: Pyralidae), 481.  
A new genus and species of Odontiinae, with remarks on *Balaenifrons* Hampson and *Taurometopa* Meyrick (Lepidoptera: Pyralidae), 865.  
New Indo-Malayan Odontiinae (Lepidoptera: Pyralidae), with taxonomic and synonymic notes, 1341.
- Nebeker, T.E. A partial life table for the Douglas-fir cone moth, *Barbara colfaxiana* (Lepidoptera: Olethreutidae), 943.  
Consumption of various Douglas-fir structures by *Barbara colfaxiana* (Lepidoptera: Olethreutidae), 1293.
- Nelson, D.L. and Baldwin, W.F. The use of radioactive platinum-iridium (IR-192) tags for locating honey bee queens in colonies (Hymenoptera: Apidae), 319.
- Neri, R.P. See Swain, W.R., 1.
- Ng, Y.S., Latheef, M.A., and Pass, B.C. Estimation of egg populations of the clover root curculio, *Sitona hispidula* (Coleoptera: Curculionidae), in Kentucky alfalfa and red clover fields, 1257.
- Nolan, R.A. See Dunphy, G.B., 341, 347.



- Novak, J.A., Akre, R.D., and Garnett, W.B. Keys to adults and puparia of five species of *Microdon* (Diptera: Syrphidae) from eastern Washington and northern Idaho, with descriptions of a new species, 663.
- Ohmart, C.P. and Dahlsten, D.L. Biological studies of bud mining sawflies, *Pleroneura* spp. (Hymenoptera: Xyelidae), on white fir in the Central Sierra Nevada of California. I. Life cycles, niche utilization, and interaction between larval feeding and tree growth, 1001.
- Okubo, A., Chiang, H.C., and Ebbesmeyer, C.C. Acceleration field of individual midges, *Anarete pritchardi* (Diptera: Cecidomyiidae), within a swarm, 149.
- O'Neal, J. See Brewer, J.W., 1153.
- Oster, G. See Wang, Y., 1359.
- Otvos, I.S. See Dunphy, G.B., 341.
- Parker, D.J. The biology of the tree-holes of Point Pelee National Park, Ontario. II. First record of *Toxorhynchites rutilus septentrionalis* in Canada (Diptera: Culicidae), 93.
- Pass, B.C. See Latheef, M.A., 1057; Ng, Y.S., 1257.
- Peck, S.B. An unusual sense receptor in internal antennal vesicles of *Ptomaphagus* (Coleoptera: Leiodidae), 81.
- Pelchat, B. See Pritchard, G., 1563.
- Pendrel, B.A. See Plowright, R.C., 967.
- Penrith, M.L. See Hamilton, W.J., 701.
- Perron, J.-M. See Perusse, F., 267; Boiteau, G., 269.
- Perusse, F., Duval, A., Chawla, S.S., and Perron, J.-M. Effects of *Taxus canadensis* on two insects (Homoptera: Aphididae; Coleoptera: Tenebrionidae), 267.
- Peschken, D.P. Host specificity of *Tingis ampliata* (Tingidae: Heteroptera): A candidate for the biological control of Canada thistle (*Cirsium arvense*), 669.
- Peterman, R.M. An evaluation of the fungal inoculation method of determining the resistance of lodgepole pine to mountain pine beetle (Coleoptera: Scolytidae) attacks, 443.
- Peters, F.E. See Madsen, H.F., 171.
- Peterson, B.V. The black flies of Iceland (Diptera: Simuliidae), 449.
- Philogène, B.J.R. and Massalski, A. Caractéristiques ultrastructurales d'une microsporidie infectant *Tyria jacobaeae* (Lepidoptère: Arctiidae), 143.
- Pierce, J. See Daterman, G.E., 875.
- Pinto, J.D. Sexual behavior in the blister beetle genus *Cysteodemus* (Coleoptera: Meloidae), 389.
- Plowright, R.C. and Pendrel, B.A. Larval growth in bumble bees (Hymenoptera: Apidae), 967.
- Porter, G.S. See Swain, W.R., 1.
- Potter, S.A. See Madsen, H.F., 171.
- Pritchard, G. See Enfield, M.A., 1425, 1435.
- Pritchard, G. and Pelchat, B. Larval growth and development of *Argia vivida* (Odonata: Coenagrionidae) in warm sulphur pools at Banff, Alberta, 1563.
- Prokopy, R.J. Attraction of *Rhagoletis* flies (Diptera: Tephritidae), to red spheres of different sizes, 593.
- Proverbs, M.D. See Jaques, R.P., 1079.
- Pulley, P.E. See Coulson, R.N., 1071.
- Pulley, P.E., Foltz, J.L., Coulson, R.N., and Martin, W.C. Evaluation of procedures for estimating within-spot populations of attacking adult *Dendroctonus frontalis* (Coleoptera: Scolytidae), 1325.
- Pulley, P.E., Foltz, J.L., Mayyasi, A.M., Coulson, R.N., and Martin, W.C. Sampling procedures for within-tree attacking adult populations of the southern pine beetle, *Dendroctonus frontalis* (Coleoptera: Scolytidae), 39.
- Quednau, F.W. See Guevrémont, H.C., 1417, 1545.
- Ramalingam, S. and Craig, G.B. The effects of JH mimic and cauterization of the corpus allatum complex on the male accessory glands of *Aedes aegypti* (Diptera: Culicidae), 897.
- Randell, R.L. See Mukerji, M.K., 229.
- Randolph, J.C. See Barlow, C.A., 1491.
- Randolph, P.A. See Barlow, C.A., 1491.
- Raske, A.G. and Bryant, D.G. Distribution, survival, and intra-tree movement of late-instar birch casebearer larvae on white birch (Lepidoptera: Coleophoridae), 1297.
- Relation of birch casebearer (Lepidoptera: Coleophoridae) numbers to per cent defoliation of white birch, 1307.
- Reid, J.A.K. and McEwen, F.L. Genetic insect control: Chromosome rearrangements isolated using a simple system in the onion maggot, *Hyalemya antiqua* (Diptera: Anthomyiidae), 1287.
- Reisen, W.K. and Emory, R.W. Intraspecific competition in *Anopheles stephensi* (Diptera: Culicidae): II. The effects of more crowded densities and the addition of antibiotics, 1475.
- The effects of larval intraspecific competition on imaginal densities in *Anopheles stephensi* (Diptera: Culicidae): A laboratory evaluation, 1481.
- Reissig, W.H. Response of the apple maggot, *Rhagoletis pomonella*, and the cherry fruit fly, *R. fausta* (Diptera: Tephritidae), to protein hydrolysate bait sprays, 161.
- Capture of released apple maggot flies, *Rhagoletis pomonella* (Diptera: Tephritidae), on sticky panels, 631.



- Retnakaran, A.** See Granett, J., 893.
- Rice, R.E. and Jones, R.A.** Monitoring flight patterns of male San Jose scale (Homoptera: Diaspididae), 1403.
- Richards, K.W.** Ovarian development of queen and worker bumble bees (Hymenoptera: Apidae) in southern Alberta, 109.
- Richards, K.W. and Richards, L.A.** A new species of *Macrocheles* (Acarina: Macrochelidae) found in bumble bee nests (Hymenoptera: Apidae), 711.
- Richards, L.A.** See Richards, K.W., 711.
- Robbins, R.G.** See Daterman, G.E., 875.
- Robert, S. and Bergeron, J.-M.** Les Siphonaptères de la région de Sherbrooke, Québec, 1571.
- Robinson, A.G.** See Rojanavongse, V., 649.
- Rojanavongse, V. and Robinson, A.G.** Species of *Aphis* (Homoptera: Aphididae) in Manitoba, with a key, and descriptions of new species, 649.
- Röling, M.P. and Kearby, W.H.** Influence of tree diameter, aspect, and month killed on the behaviour of scolytids infesting black oaks, 1235.
- Rosenberg, D.M.** See Madder, M.C.A., 891.
- Rowell, H.J.** See Chant, D.A., 1605.
- Safranyik, L.** See Dyer, E.D.A., 77.
- Salkeld, E.H.** Biosystematics of the genus *Euxoa* (Lepidoptera: Noctuidae). IX. Morphological and electrophoretic characteristics of the eggs of the "declarata group", 1137.
- Sanders, C.J., Daterman, G.E., and Ennis, T.J.** Sex pheromone responses of *Choristoneura* spp. and their hybrids (Lepidoptera: Tortricidae), 1203.
- Sanford, K.H.** See Jaques, R.P., 1079.
- Schenk, J.A.** See Kirtibutr, N., 381.
- Schlinger, E.I.** See Dahlsten, D.L., 727.
- Schmid, J.M.** See Frye, R.H., 1221.
- Schmidt, F.H.** Differences in thermal requirements for diapause termination in two western *Choristoneura* spp. (Lepidoptera: Tortricidae), 1469.
- Scudder, G.G.E.** The World Rhyparochrominae (Hemiptera: Lygaeidae). XVI. The status of *Tethallotrum* Scudder and the description of a new species of *Alloccentrum* Bergroth from Madagascar, 537.
- Sebastien, R.J.** See Baldwin, W.F., 249.
- Singh, N.B.** See Bailey, C.G., 687.
- Sharpe, P.J.H.** See Barfield, C.S., 1503.
- Smirnov, W.A.** Confirmations expérimentales du potentiel du complexe *Bacillus thuringiensis* et chitinase pour la répression de la tordeuse des bourgeons de l'épinette, *Choristoneura fumiferana* (Lepidoptera: Tortricidae), 351.
- See also McNeil, J.N., 37.
- Smith, B.C. and Halliday, C.G.** Reduction in damage to potatoes by *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae) by mixtures of insecticides and silica, 59.
- Smith, D.S. and Holmes, N.D.** The distribution and abundance of adult grasshoppers (Acrididae) in crops in Alberta, 1918-1975, 575.
- Smith, I.M.** A new species of *Mideopsis* (*Nudomideopsis*) (Acari: Parasitengona: Mideopsidae) from North America, 533.
- Two new species of *Trisetacus* (Prostigmata: Eriophyoidea) associated with Cupressaceae, 843.
- Descriptions of larvae of some axonopsine water mites (Prostigmata: Parasitengona: Aturidae), with notes on phylogenetic affinities of the subfamily, 953.
- A new species of eriophyoid mite with eye-like structures, and remarks on the genus *Phytoptus* (Acari: Prostigmata: Phytoptidae), 1097.
- Smith, J.J.B.** See Friend, W.G., 15.
- Smith, J.J.B. and Friend, W.G.** A serial sectioning technique for morphological study of complex chitinous structures with the scanning electron microscope, 477.
- Smith, L.B.** Efficiency of Berlese-Tullgren funnels for removal of the rusty grain beetle, *Cryptolestes ferrugineus*, from wheat samples, 503.
- Solomon, J.D.** Frass characteristics for identifying insect borers (Lepidoptera: Cossidae and Sesiidae; Coleoptera: Cerambycidae) in living hardwoods, 295.
- Soutar, B.M.** See Gibo, D.L., 615.
- Steck, W., Underhill, E.W., Chisholm, M.D., Bailey, B.K., Loeffler, J., and Devlin, C.G.** Sex attractants for males of 12 moth species found in Western Canada, 157.
- Steck, W.F.** See Underhill, E.W., 1335; Struble, D.L., 1393.
- Stevens, R.E., Borg, T.K., and Thatcher, T.O.** Notes on a pine-feeding budworm, *Choristoneura lambertiana ponderosana* (Lepidoptera: Tortricidae), in the Colorado Rockies, 1269.
- Stinner, R.E., Jones, J.W., Tuttle, C., and Caron, R.E.** Population mortality and cyclicity as affected by intraspecific competition, 879.

- Struble, D.L. Bioassay of sex pheromones in the army cutworm, *Euxoa auxiliaris*; the pale western cutworm, *Agrotis orthogonia*; and the *Euxoa tristicula* (Lepidoptera: Noctuidae), 365.
- Struble, D.L. and Lilly, C.E. An attractant for the beet webworm, *Loxostege sticticalis* (Lepidoptera: Pyralidae), 261.
- Struble, D.L. and Swales, G.E. Sex attractant for clover cutworm, *Scotogramma trifolii*: Field tests with various ratios and Z-11-hexadecen-1-yl acetate and Z-11-hexadecen-1-ol, and with various quantities of attractant on two types of carriers, 369.
- Struble, D.L., Swales, G.E., and Ayre, G.L. A sex attractant for males of the darksided cutworm, *Euxoa messoria* (Lepidoptera: Noctuidae), 975.
- Struble, D.L., Swales, G.E., Steck, W.F., Underhill, E.W., and Chisholm, M.D. A sex attractant for *Leucania commoides*: A mixture of Z-9-tetradecen-1-yl acetate, Z-11-hexadecen-1-yl acetate, and Z-11-hexadecen-1-ol, 1393.
- Stuart, R.J. See Baldwin, W.F., 249.
- Sullivan, C.R., Griffiths, K.J., and Wallace, D.R. Low winter temperatures and the potential for establishment of the egg parasite *Anastatus disparis* (Hymenoptera: Eupelmidae) in Ontario populations of the gypsy moth, 215.
- Summers, C.G. See Christensen, J.B., 1599.
- Svec, J.H. See Bond, E.J., 285.
- Swales, G.E. See Struble, D.L., 369, 975, 1393.
- Swain, W.R., Wilson, R.M., Neri, R.P., and Porter, G.S. A new technique for remote monitoring of activity of freshwater invertebrates with special reference to oxygen consumption by naiads of *Anax* sp. and *Somatochlora* sp. (Odonata), 1.
- Syme, P.D. Observations on the longevity and fecundity of *Orgilus obscurator* (Hymenoptera: Braconidae) and the effects of certain foods on longevity, 995.
- Temporale, A. See Gibo, D.L., 615.
- Teskey, H.J. and Chilcott, J.G. A revision of the Nearctic species of *Syndyas* Loew (Diptera: Empididae), 1445.
- Teskey, H.J. and Valiela, I. The mature larva and puparium of *Canace macateei* (Diptera: Canaceidae), 545.
- Thatcher, T.O. See Stevens, R.E., 1269.
- Thomson, A.J. and Holling, C.S. A model of carbohydrate nutrition in the blowfly *Phormia regina* (Diptera: Calliphoridae), 1181.
- Tomlin, A.D. Toxicity of soil applications of the fungicide benomyl and two analogues to three species of Collembola, 1619.
- Turnbull, S.A. See Harris, C.R., 1109.
- Tuskes, P.M. Observations on the biology of *Parargyractis confusalis*, an aquatic pyralid (Lepidoptera: Pyralidae), 695.
- Tuttle, C. See Stinner, R.E., 879.
- Underhill, E.W. See Steck, W., 157; Struble, D.L., 1393.
- Underhill, E.W., Steck, W.F., and Chisholm, M.D. A sex pheromone mixture for the bertha armyworm moth, *Mamestra configurata*: (Z)-9-tetradecen-1-ol acetate and (Z)-11-hexadecen-1-ol acetate, 1335.
- Urquhart, F.A. and Urquhart, N.R. Overwintering areas and migratory routes of the monarch butterfly (*Danaus p. plexippus*, Lepidoptera: Danaidae) in North America, with special reference to the western population, 1583.
- Urquhart, N.R. See Urquhart, F.A., 1583.
- Valiela, I. See Teskey, H.J., 545.
- van den Driessche, P. See Barclay, H., 763.
- Veazey, J.N. See Kay, C.A.R., 721.
- Wallace, D.R. See Sullivan, C.R., 215.
- Wang, Y. See Gutierrez, A.P., 1375, 1457.
- Wang, Y., Gutierrez, A.P., Oster, G., and Daxl, R. A population model for plant growth and development: Coupling cotton-herbivore interaction, 1359.
- Weber, B.C. Parasitoids of the introduced pine sawfly, *Diprion similis* (Hymenoptera: Diprionidae), in Minnesota, 359.
- Wellington, W.G. See Mackay, P.A., 53.
- Wenz, J.M. See Dahlsten, D.L., 727.
- Werner, R.A. Diapause termination in overwintering pupae of *Rheumaptera hastata* (Lepidoptera: Geometridae) in interior Alaska, 1149.
- West, A.S. See Baldwin, W.F., 249.
- Westphal, D. See Gutierrez, A.P., 1457.
- Wheeler, A.G. Studies on the arthropod fauna of alfalfa. VII. Predaceous insects, 423.
- Whitcomb, W.H. See Kay, C.A.R., 721.
- White, N.D. and Laing, J.E. Some aspects of the biology and a laboratory life table of the acarine predator *Zetzellia mali*, 1275.

- Wiens, A.P.** See Madder, M.C.A., 891.
- Wilson, G.G.** Effects of the microsporidia *Nosema disstriae* and *Pleistophora schubergi* on the survival of the forest tent caterpillar, *Malacosoma disstria* (Lepidoptera: Lasiocampidae), 1021.
- Wilson, M.L.** See Baldwin, W.F., 249.
- Wilson, R.M.** See Swain, W.R., 1.
- Wong, H.R.** Chinese species of *Pristiphora* and their relationship to Palaearctic and Nearctic species (Hymenoptera: Tenthredinidae), 101.  
*Fallocampus*: A new sawfly genus for the Nearctic species of *Platycampus* Schiödte (Hymenoptera: Tenthredinidae), 1103.
- Wylie, H.G.** Observations on *Athrycia cinerea* (Diptera: Tachinidae), a parasite of *Mamestra configurata* (Lepidoptera: Noctuidae), 747.  
Observations on *Mericia ampelus* (Diptera: Tachinidae), an occasional parasite of bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae), in Western Canada, 1023.  
Preventing and terminating pupal diapause in *Athrycia cinerea* (Diptera: Tachinidae), 1083.
- Wylie, H.G. and Bracken, G.K.** Observations on *Ichneumon canadensis*, a parasite of noctuid pupae in southern Manitoba, 1295.
- Wylie, H.G. and Bucher, G.E.** The bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae). Mortality of immature stages on the rape crop, 1972-1975, 823.
- Yarger, L.C. and Brewer, J.W.** Biology and habits of the white fir needle miner, *Epinotia meritana* (Lepidoptera: Olethreutidae), in Colorado, 849.
- Yeargan K.V.** See Latheef, M.A., 1057.
- Ying-Shin Peng and Jay, S.C.** Larval rearing by worker honey bees lacking their mandibular glands. I. Rearing by small numbers of worker bees, 1175.
- Yoshimoto, C.M.** A new species of *Spalangiopecta* Masi in North America (Chalcidoidea: Pteromalidae, Ceinae), 541.  
The North American species of the genus *Achrysocharoides* (Hymenoptera: Eulophidae), 907.  
A new species of *Ooencyrtus leptoglossi* (Hymenoptera: Chalcidoidea, Encyrtidae) reared from eggs of *Leptoglossus corculus* (Hemiptera: Coreidae), 1009.  
Revision of the Diparinæ (Pteromalidae: Chalcidoidea) from America north of Mexico, 1035.
- Young, J.W.** See Furniss, M.M., 1063.

# THE ENTOMOLOGICAL SOCIETY OF CANADA

## SUPPLEMENTS AND MEMOIRS

- No. 1 of Vol. LXXXVII (mailed 30 March 1955): The comparative internal larval anatomy of sawflies (Hymenoptera: Symphyta) — Doreen E. Maxwell
  - No. 1 of Vol. LXXXVIII (mailed 10 December 1956): Revision of the Nearctic species of *Agriotes* (Coleoptera: Elateridae) — Edward C. Becker
  - No. 2 of Vol. LXXXVIII (mailed 16 January 1957): Leafhoppers (Homoptera: Cicadellidae) of Canada and Alaska — Bryan P. Beirne
  - No. 3 of Vol. LXXXVIII (mailed 2 January 1957): The New World species of *Chrysomela* L. (Coleoptera: Chrysomelidae) — W. J. Brown
- NOTE: After this issue the "Supplements" became, in fact, a self-contained series with a continuous numbering system independent of *The Canadian Entomologist*. Nevertheless, they were still issued (with later exceptions) as "Accompanying" volumes of *The Canadian Entomologist*.  
Because four Supplements had already been published, the next one was called No. 5 — there is no Supplement No. 4.
- No. 4 — there is no Supplement No. 4.
  - No. 5 (mailed 2 December 1957): The use of larval anatomy in the study of bark beetles (Coleoptera: Scolytidae) — J. B. Thomas
  - No. 6 (mailed 8 April 1958): Taxonomy, life history, and habits of the elliptoid-eyed species of *Schinia* (Lepidoptera: Noctuidae), with notes on the Heliethidinae — David F. Harwick
  - No. 7 (mailed 28 April 1958): The Archipinae of North America (Lepidoptera: Tortricidae) — T. N. Freeman
  - No. 8 (mailed 12 September 1958): Larvae of the Nearctic Laurentiinae (Lepidoptera: Geometridae) — W. C. McGuffin
  - No. 9 (mailed 14 November 1958): Acridoidea of southern Alberta, Saskatchewan, and Manitoba (Orthoptera) — Arthur R. Brooks
  - No. 10 (mailed 26 March 1959): Larvae of the North American Olethreutidae (Lepidoptera) — Margaret Rae MacKay
  - No. 11 (mailed 26 March 1959): Male genitalia as taxonomic characters in the Miridae (Hemiptera) — Leonard A. Kelton
  - No. 12 (mailed 15 June 1959): Phytoseiid mites (Acarina: Phytoseiidae). I. Bionomics of seven species in southeastern England. II. A taxonomic review of the family Phytoseiidae, with descriptions of 38 new species — D. A. Chant
  - No. 13 (mailed 14 November 1960): A synopsis of the genus *Rhopalosiphum* in Canada (Homoptera: Aphididae) — W. R. Richards
  - No. 14 (mailed 16 March 1961): A revision of the Nearctic species of Fanniinae (Diptera: Muscidae) — J. G. Chillcott
  - No. 15 (mailed 1 February 1961): Synopsis of Nearctic Ichneumoninae Stenopneusticae with particular reference to the northeastern region (Hymenoptera). I. Introduction, key to Nearctic genera of Ichneumoninae Stenopneusticae, and synopsis of the Protichneumonini north of Mexico — Gerd H. Heinrich
  - No. 16 (mailed 11 January 1961): Needle-mining Lepidoptera of pine in North America — T. N. Freeman
  - No. 17 (mailed 25 October 1961): The classification of the Papilionidae (Lepidoptera) — Eugene Munroe
  - No. 18 (mailed 4 April 1961): Synopsis of Nearctic Ichneumoninae Stenopneusticae with particular reference to the northeastern region (Hymenoptera). II. Synopsis of the Ichneumonini: Genera *Orgichneumon*, *Cratichneumon*, *Homotherus*, *Aculichneumon*, *Spilichneumon* — Gerd H. Heinrich
  - No. 19 (mailed 4 April 1961): Mathematical models for use in insect pest control — Kenneth E. F. Watt
  - No. 20 (mailed 6 April 1961): Adult Elateridae of southern Alberta, Saskatchewan and Manitoba (Coleoptera) — A. R. Brooks
  - No. 21 (mailed 13 September 1961): Synopsis of Nearctic Ichneumoninae Stenopneusticae with particular reference to the northeastern region (Hymenoptera). III. Synopsis of the Ichneumonini: Genera *Ichneumon* and *Thyrateles* — Gerd H. Heinrich
  - No. 22 (mailed 29 May 1961): Taxonomy and distribution of Nearctic *Vespula* — C. D. F. Miller

